

## Isolation and Characterization of Estra-2<sup>II</sup>-en -17-ol, 3yl benzoate from *Mucuna pruriens* ( *Utilis*).

Rosemary Izunwanne Uchegbu <sup>1\*</sup> and JohnBull Onyekachi Echeme <sup>2</sup>

\*1. Department of Chemistry, Alvan Ikoku Federal College of Education, P. O. Box 1033, Owerri. Imo State. Nigeria.

2. Department of Chemistry, Michael Okpara University of Agriculture Umudike. Umuahia . Abia State . Nigeria.

\*Corresponding author, [mrsriu@yahoo.com](mailto:mrsriu@yahoo.com).

### Abstract

Chemical investigation of the bioactive constituents of the seeds of *Mucuna pruriens* (*utilis*) resulted in the isolation of a new steroid, Estra-2<sup>II</sup>-en -17-ol, 3yl benzoate. The structure was elucidated using NMR spectroscopy in combination with IR and MS spectra data.

**Keywords:** *Mucuna*, anti-microbial, anti-oxidant, anti-inflammatory, anti-hypertensive.

### 1. Introduction

Plants have been known to be useful to man; as source of food and drug. They are of paramount importance in the treatment of diseases for man. Nigeria is blessed with many of such plants and they are used as food and in herbal medicine to cure diseases and heal injuries. Some of these plants exhibit anti-microbial, anti-oxidant, anti-inflammatory and anti-hypertensive activities. Such medicinal plants include *Mucuna pruriens* (*utilis*). *M. pruriens* is extensively used in herbal medicine and as food. The seeds have been found to contain high levels of protein and carbohydrate and thus the flour has been recommended to be incorporated into cereals and root/tuber food products to improve their nutritional quality (Udensi and Eke, 2000). The flour is used in soup as soup thickener and to enhance the flavour of the soup. *M. pruriens* is grown widely as a green vegetable, both the green pods and the mature beans are boiled and eaten as food (Burkill, (1995); Duke, (1981)). The seeds are also roasted and ground to make a coffee substitute. Extracts from the seeds have been recommended as one of the best tonics to improve vitality (Manyam et al, 2004).

*M. pruriens* is highly valued because of its medicinal use. It is used in herbal medicine as a powerful nervine tonic and an effective aphrodisiac in the treatment of disorders of the male and female reproductive systems (Numida, 2009; Ezueh, 1997; Amin et al, 1996). The seeds are effective in the treatment of parkinson diseases (Manyam et al, 2004; Katzenschlager et al, 2004). Extracts from the seeds have been used to enhance sexual stamina, sensation and libidos in both men and women due to its dopamine inducing properties (Amin et al, 1996; Giuliano and Allard, 2001).

*Mucuna pruriens* seeds have also been found to have antidepressant properties when consumed. It is a very beneficial supplement for body builders as it increases the body's ability to build lean muscle, breakdown fat, and is highly recommended to sportsmen (Ray Sahelian, M.D).

From literature, it is obvious that *Mucuna pruriens* seeds contain some hormones since it is used to build lean muscles and as aphrodisiac by the natives, but isolation has not been made to that effect, thus this study was undertaken to isolate and characterize bioactive constituents of *Mucuna pruriens* seeds.

### 2. Materials and Method.

**2.1 Plant Materials :** The seeds of *M. pruriens* were harvested from the field of National Root Crops Research Institute Umudike ,Abia state, Nigeria. Authentication of plant materials was done by Dr A. Nmergini of Taxonomy section, Forestry Department, Micheal Okpara University of Agriculture, Umudike, Nigeria.


**2.2 Extraction and Isolation of Plant Materials :** The seeds were cracked, the testa were removed and the seeds were milled into fine powder with Thomas Willey milling machine and then stored in air tight bottles for analysis. 2kg of the sample was percolated in 98% ethanol for 48hrs, this was then filtered. The filtrate was concentrated with Rotary evaporator at 40°C to a dark brown crude extract (44.3g). The crude extract was partitioned between CHCl<sub>3</sub> and water and a CHCl<sub>3</sub> - soluble fraction (17.5g) was obtained. 10.0g of the CHCl<sub>3</sub> fraction was then partitioned between petroleum ether (60 – 80°C) and aqueous methanol. 5.0g of the CHCl<sub>3</sub> fraction was then subjected to column chromatography over silica gel (200 mesh) and eluted gradually with 100ml petroleum ether, then petroleum ether : CHCl<sub>3</sub> (90:10; 80:20; 70:30; 60:40; 50:50; 40:60; 30:70; 20:80; 10:90;), and 100ml CHCl<sub>3</sub>; then CHCl<sub>3</sub> : Methanol (90:10; 80:20; 70:30; 60:40; 50:50; 40:60; 30:70; 20:80; 10:90) and 100ml methanol to yield ten major fractions labeled : G (yellow oil, 0.3g), I ( yellow oil, 0.4g), b (brown oil, 0.3g) c<sub>1</sub> (yellow oil, 0.6g), sc<sub>2</sub> (yellow oil, 0.60g), d (brown oil 0.32g) f<sub>2</sub> (brown solid, 0.3g), g<sub>2</sub> (brown oil, 0.5g), h (dark brown oil, 0.4g), i (light yellow oil, 0.3g) and i<sub>2</sub> (light yellow oil, 0.4g).

Chromatographic (partition chromatography, column chromatography, and TLC) and spectroscopic (IR,  $^1\text{H}$ NMR,  $^{13}\text{C}$ NMR and MS) techniques were employed to isolate, characterize and identify active constituents from  $\text{CHCl}_3$  extracts of the seeds.

### 3.0 Results and Discussions:

Compound [1] was eluted with methanol and chloroform at the ratio of 70:30. It was isolated as yellow oil (0.6g). Based on combined analysis of HREIMS ( $m/z$  445.41),  $^1\text{H}$ NMR,  $^{13}\text{C}$ NMR and IR spectra, compound [1] was proposed as Estra-2''-en-17-ol, 3yl benzoate with the molecular formula  $\text{C}_{29}\text{H}_{34}\text{O}_4$ ,  $m/z$  446 calculated.

Analysis of the IR is shown in Table 1 and the analysis of  $^1\text{H}$ NMR and  $^{13}\text{C}$ NMR are shown in Table 2. The IR spectrum revealed the presence of hydroxyl, aliphatic, carbonyl, aromatic and ether bands at ( $3470.52\text{cm}^{-1}$ ,  $2924.97\text{cm}^{-1}$ ,  $1741.0\text{cm}^{-1}$ ,  $3008.59\text{cm}^{-1}$  and  $1169.89\text{cm}^{-1}$ ) respectively.



[1] Estra-2''-en-17-ol, 3yl benzoate

**Table 1: Infra-red Analysis of Estra-2''-en-17-ol, 3yl benzoate [1] from *M. pruriens***

IR Absorption ( $\text{cm}^{-1}$ )	Functional Group	Compound
3470.52	OH	Alcohol
3007.89	C=C	Aromatic
2925.44	-CH <sub>2</sub>	Aliphatic
2855.35	-CH <sub>2</sub>	Aliphatic
1742.73	C=O	Carbonyl
1167.76	C-O	Ether

**Table 2:  $^1\text{H}$  and  $^{13}\text{C}$ NMR Analysis of Estradiol-17-ol-3yl benzoate.**

	$\delta\text{C}$			$\delta\text{H}$	
1	127.95	CH	5.323	IHd	CH
2	128.12	CH	5.342	IHd	CH
3	130.06	C			
4	129.76	CH	5.350	IHs	CH
5	130.27	C			
6	31.59	CH <sub>2</sub>	1.231	2Ht	CH <sub>2</sub>
7	31.72	CH <sub>2</sub>	1.254	2Ht	CH <sub>2</sub>
8	33.66	CH	1.607	IHd	CH
9	33.99	CH	1.629	IHd	CH
10	129.76	C	—	—	—
11	31.75	CH <sub>2</sub>	1.278	2Hq	CH <sub>2</sub>
12	31.84	CH <sub>2</sub>	1.285	2Ht	CH <sub>2</sub>
13	34.11	C	—	—	—
14	34.14	CH	1.376	IHt	CH
15	31.99	CH <sub>2</sub>	1.302	2Ht	CH <sub>2</sub>
16	34.75	CH	2.080	IHd	CH
17	77.11	C—OH	7.265	IHs	OH
18	14.13	CH <sub>3</sub>	0.857	3Hs	CH <sub>3</sub>
1'	127.95	C			
2'	128.12	CH	5.360	IHd	CH
3'	130.06	CH	5.370	IHt	CH
4'	129.76	CH	5.323	IHt	CH
5'	130.27	CH	5.342	IHt	CH
6'	129.76	CH	5.350	IHd	CH
7'	173.38	C=O			
1''	76.69	CH <sub>2</sub>	5.386	2Ht	CH <sub>2</sub>
2''	172.92	CH	2.347	IHq	CH
3''	173.34	CH	2.373	IHm	CH
4''	14.18	CH <sub>3</sub>	0.857	3Hd	CH <sub>3</sub>

**S = singlet, d = doublet, t = triplet, q = quartet, m = multiplet**

The mass spectrum of compound [1] showed  $m/z$  445.41 which was calculated for  $\text{C}_{29}\text{H}_{34}\text{O}_4$  ( $m/z$  446) and base peak  $m/z$  55.18 calculated for  $\text{C}_4\text{H}_7$  ( $m/z$  55). The  $^1\text{H}$ NMR spectrum of compound [1] showed presence of methylene protons observed at  $\delta\text{H}$  1.231,  $\delta\text{H}$  1.254,  $\delta\text{H}$  1.278,  $\delta\text{H}$  1.285, methine protons at  $\delta\text{H}$  1.607 -  $\delta\text{H}$  2.373 and aromatic protons which showed their signals at  $\delta\text{H}$  5.232 -  $\delta\text{H}$  5.370. The spectrum indicated presence of hydroxyl proton at  $\delta\text{H}$  7.265.

The  $^{13}\text{C}$ NMR spectrum depicted the presence of aromatic carbon atoms, their chemical shifts were observed at  $\delta$  127.95 -  $\delta$  130.27. The carbonyl chemical shifts appeared at  $\delta$  173.38. The methyl carbons showed their signals at  $\delta$  14.13 and  $\delta$  14.18. The signals of the methylene carbons were observed at  $\delta$  31.59 -  $\delta$  31.99 while the chemical shifts of the methine carbons appeared at  $\delta$  33.66 -  $\delta$  34.25.

Fragmentation pattern of compound [1] is shown in fig (1). Detachment of  $\text{C}_4\text{H}_7$  ion from the compound afforded the base peak  $m/z$  55.18 calculated for  $m/z$  55. The cleavage between the two oxygen atoms in the benzoate gave the peak at  $m/z$  105. Also detachment of  $\text{C}_5\text{H}_5\text{O}$  ion from the compound afforded another prominent peak  $m/z$  81.20 calculated for  $m/z$  81. The isolated compound has the skeleton of Estradiol, a sex hormone. The proton in the second hydroxyl is replaced by phenyl oxide. Estradiol protects against brain injury, neurodegeneration and cognitive decline. It protects against stroke injury (Dubal et al, 2000). Thus, this compound isolated from *M. pruriens* has biological and physiological activities. The presence of this compound in *Mucuna pruriens* seed may be the reason why the natives use this plant in the treatment of disorders of the male and female reproductive systems, parkinson disease and increase libido in both men and women (Numida, 2009; Ezueh, 1997; Amin et al, 1996). Estradiol has been found to reduce anxiety and depression - like behavior of aged female mice (Walf and Frye, 2005). The isolated compound may also be the reason *mucuna pruriens* seeds are used by the herbalists as anti-depressant.

## Conclusion

The results presented here showed that *M. pruriens* plant can serve as a pharmacodynamic agent. The plant can be seen as a potential source of useful drugs and can serve as source of quality raw material for pharmaceutical

industries. Thus this contributes to the scientific evidence for the use of this medicinal plant in traditional medicine for the treatment of diseases like pains, depressions, erectile dysfunction, parkinson disease, e.t.c.




Fig 1: fragmentation Pattern of Compound [1].

#### Acknowledgement

We are grateful to Dr Nkechi Ibisi of Chemistry department, Micheal Okpara University of Agriculture, Umudike, Nigeria for running our spectra in China and to Dr A. Nmeregini of Taxonomy section, Forestry department, Micheal Okpara University of Agriculture, Umudike, Nigeria for authenticating our sample.

#### References

- Amin, K.M.Y, Khan, M.N and Zillur-Rehman S, (1996) "Sexual function improving effect of *Mucuna pruriens* in sexually normal male rats".
- Burkill, H.M (1995). The useful Plants of West Tropical Africa. (vol 3). Royal Botanic Gardens, London .Pp 101.
- Duke J.A. (1981). Handbook of legumes of world economic importance. Plenum press. New York.
- Dubal, D.B, Zhu, H. Yu, J., Rau, S.W., Shughrue, P.J., Merchenthaler, I., Kindy, M.S. and Wise, P.M., (2000). "Estrogen receptor  $\alpha$ , not  $\beta$ , is a critical link in estradiol-mediated protection against brain injury". *Proc. Natl. Acad. Sci. USA*, 10, 1073.
- Ezueh, M.I (1997) Cultivation and Utilization of Minor Food Legumes in Nigeria. Tropical Grain legume Information Center, IITA, Ibadan, Nigeria
- Giuliano, F. and Allard, J. (2001). Dopamine and sexual function. *Int J Impot Res* 13 Suppl 3:18-28
- Katzenschlager R, Evans A, and Manson, A (2004). *Mucuna pruriens* in Parkinson's disease: a double blind clinical and pharmacological study. *J Neurol Neurosurg Psychiatry* 75:1672-1677
- Manyam, B.U., Dhanasekaran, M. and Hare, T.A (2004). Neuroprotective effects of the antiparkinson drug *Mucuna Pruriens*. *Phytother Res* 18:97-101.
- Numida, M.L (2009). "The effects of Raw and Processed *Mucuna pruriens* seed Based Diets on the growth parameters and meat characteristics of Benin Local Guinea Fowl". *International journal of Poultry sciences* 8(9).
- Ray Sahelian, M.D. Accessed at [www.raysahelian.com/mucunapruriens.html](http://www.raysahelian.com/mucunapruriens.html).
- Udensi, E.A and Eke, O. (2000). Proximate Composition and Functional Properties of Flour Produced from *Mucuna cochinchinensis* and *Mucuna utilis*. Food and Fiber production in Nigeria in the 21<sup>st</sup> century. Pg 170 – 173.
- Walf, A.A and Frye, C.A. (2005). Anxiety and anti-depressive behavior produced by Physiological estradiol regimen may be modulated by hypothalamic – pituitary adrenal axis activity. *Neuropsychopharmacology* 30(7): 1288 – 1301.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

## CALL FOR JOURNAL PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> The IISTE editorial team promises to the review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

## MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Recent conferences: <http://www.iiste.org/conference/>

## IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

